

Mr Rhys Thomas FRCS
Consultant General and
Upper Gastrointestinal Surgeon

Shirley Oaks Hospital,
Poppy Lane,
Shirley Oaks Village,
Croydon,
CR9 8AB.

Patient
Information
for Consent

GS03lite Femoral Hernia Repair

Expires end of May 2017
Issued April 2016

Local information

If you would like more information or if you would like to contact us please use any of the following:

Tel: 0208 655 5500
E-mail: rlt197@doctors.org.uk
Website: www.croydonsurgeon.com
Twitter: [@croydonsurgeon](https://twitter.com/croydonsurgeon)

Get more information and references at www.aboutmyhealth.org
Tell us how useful you found this document at www.patientfeedback.org



www.rcseng.ac.uk

www.bads.co.uk

www.rcsed.ac.uk

www.asgbi.org.uk

www.pre-op.org



This document will give you information about a femoral hernia repair. If you have any questions, you should ask your GP or other relevant health professional.

What is a femoral hernia?

Weak spots can develop in the layer of muscle in your abdominal wall, resulting in the contents of your abdomen pushing through. This produces a lump called a hernia (see figure 1).




Figure 1
A femoral hernia

A femoral hernia happens at the hole in the wall of your abdomen where the femoral artery and vein pass from your abdomen into your leg. A hernia can be dangerous because your intestines or other structures within your abdomen can get trapped and have their blood supply cut off (strangulated hernia).

What are the benefits of surgery?

You should no longer have the hernia. Surgery should prevent the serious complications that a hernia can cause.

Are there any alternatives to surgery?

The hernia will not get better without surgery.

What does the operation involve?

Various anaesthetic techniques are possible. The operation usually takes about 45 minutes.

Your surgeon will make a cut either directly above the lump or a little higher up and will remove the 'hernial sac'. They will narrow the hole (femoral canal) through which the contents of the abdomen passed, using stitches or a synthetic mesh. Your surgeon will close your skin.

What complications can happen?

1 General complications

- Pain
- Bleeding
- Infection of the surgical site (wound)
- Unsightly scarring
- Blood clots

2 Specific complications

- Developing a lump under your wound
- Difficulty passing urine
- Injury or narrowing of the femoral vein
- Injury to structures that come from your abdomen and are within the hernia
- Temporary weakness of your leg
- Injury to nerves

How soon will I recover?

You should be able to go home the same day. Increase how much you walk around over the first few days.

You should be able to return to work after two to four weeks, depending on the extent of surgery and your type of work.

Regular exercise should help you to return to normal activities as soon as possible. Before you start exercising, ask the healthcare team or your GP for advice.

The hernia can come back.

Summary

A femoral hernia is a common condition caused by a weakness in your abdominal wall, near the femoral canal. If left untreated, a femoral hernia can cause serious complications.

Acknowledgements

Author: Prof Simon Parsons DM FRCS (Gen. Surg.)
Illustrations: Medical Illustration Copyright ©
Medical-Artist.com

This document is intended for information purposes only and should not replace advice that your relevant health professional would give you.